


CS 428: Fall 2010

# Introduction to Computer Graphics

Viewing and  
projective transformations

# Modeling and viewing transformations



# Modeling and viewing transformations

- OpenGL order

```
glMatrixMode (GL_MODELVIEW)
```

```
glLoadIdentity ()
```

```
glMultMatrix (V)
```


```
glMultMatrix (M)
```

```
draw () ← Transformation is VM
```


- In OpenGL, these transformations are place on the **modelview** matrix stack
- The **projection** matrix stack is only for storing the projection matrix resulting from **glOrtho ()**, **glFrustum ()**, or **gluProjection ()**

# 3D viewing

- The eye has a view cone
  - Approximated in CG by a “rectangular cone” = square **frustum**
  - Good for rectangular viewing window




# 3D viewing process




# Truncated view volumes

- Orthographic



# Truncated view volumes


- Perspective



on near plane  
gl Frustum( $l, r, b, t, n, f$ )  
glu Perspective( $fov-y, aspect, n, f$ )

# Where's the film?

- A rectangle with known aspect ratio on the infinite film plane
- “Where” doesn't matter, as long as the film plane is parallel to *far* and *near*
- Will be scaled to viewport coordinates





# OpenGL projection matrices

- How is this implemented in OpenGL?
- The following matrices assume
  - Camera center at origin
  - Looking down negative z-axis
  - y-axis is “up”
  - $near, far > 0$
  - $right = -left = 1$
  - $top = -bottom = 1$
- See `glOrtho()` etc. manpages for general case



# OpenGL projection matrices

- Orthographic projection

$$P_{ortho} = \begin{bmatrix} 1 & & & \\ & 1 & & \\ & & \frac{2}{n-f} & \frac{n+f}{n-f} \\ & & & 1 \end{bmatrix}$$

$$P_{ortho} \cdot \begin{bmatrix} x \\ y \\ \frac{z}{n} \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ -y \\ 1 \end{bmatrix}$$

$$P_{ortho} \cdot \begin{bmatrix} x \\ y \\ -\frac{z}{f} \\ 1 \end{bmatrix} = \begin{bmatrix} x \\ y \\ \frac{z}{f} \\ 1 \end{bmatrix}$$

maps ortho view volume to  $x, y, z \in [-1, 1]$  (cube @ origin)

where near  $\rightarrow z = -1$

far  $\rightarrow z = +1$

normalized  
device  
coordinates

# OpenGL projection matrices

- Perspective projection

$$P_{\text{persp}} = \begin{bmatrix} c/r & & & \\ & c/t & & \\ & & \frac{n+f}{n-f} & \frac{2nf}{n-f} \\ & & -1 & \end{bmatrix}$$

$$P_{\text{persp}} \cdot \begin{bmatrix} x \\ -y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} \left(\frac{n}{r}\right)x \\ \left(-\frac{n}{t}\right)y \\ -z \\ n \end{bmatrix} \xrightarrow{\div w} \begin{bmatrix} x/r \\ y/t \\ -1 \\ 1 \end{bmatrix}$$

$$P_{\text{persp}} \cdot \begin{bmatrix} x \\ y \\ -f \\ 1 \end{bmatrix} = \begin{bmatrix} \left(\frac{n}{r}\right)x \\ \left(\frac{n}{t}\right)y \\ f \\ f \end{bmatrix} \xrightarrow{\div w} \begin{bmatrix} \frac{n}{r} \cdot \frac{x}{f} \\ \frac{n}{t} \cdot \frac{y}{f} \\ 1 \\ 1 \end{bmatrix}$$

maps frustum to cube @ origin  $[-1, 1]^3$


$$P_{\text{persp}} \cdot \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{n}{r}x \\ \frac{n}{t}y \\ z \\ -z \end{bmatrix} \xrightarrow{\div w} \begin{bmatrix} -\frac{n}{r} \frac{x}{z} \\ -\frac{n}{t} \frac{y}{z} \\ 1 \\ 1 \end{bmatrix}$$

$x, y$  divided by  $z \rightarrow$  foreshortening

# Perspective projections

- Perspective projections are not affine transformations
  - Relative lengths are no longer invariant
  - Distant objects (of same size) are made smaller than near ones (= foreshortening)
  - Given a projected point  $P=(x, y)^t$  and eye position  $A=(-x_0, 0)^t$ , then by the theorem of intersecting lines the image  $B$  is  $(0, y_0)^t$

$$\frac{y_0}{y} = \frac{x_0}{x + x_0}$$



# Perspective projections

- In general, the mapping is

$$y_0 = y \cdot \frac{x_0}{x_0 + x}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 0 \\ \frac{y \cdot x_0}{x_0 + x} \end{pmatrix}$$

- Resulting in the homogeneous  $3 \times 3$ -Matrix

- 2D-Geometry!

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & x_0 & 0 \\ 1 & 0 & x_0 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{x_0} & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} 0 \\ x_0 \cdot y \\ x + x_0 \end{bmatrix} = \begin{bmatrix} 0 \\ \frac{x_0}{x + x_0} \cdot y \\ 1 \end{bmatrix}$$

# Perspective projections

- The projection is composed of two transformations
  - The perspective transformation
  - The subsequent parallel projection

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{x_0} & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{x_0} & 0 & 1 \end{bmatrix}$$

**Perspective  
projection**


**Parallel-  
projection**

**Perspective  
transformation**

# Perspective transformations


- Properties of perspective transformations of the form

$$T_p \cdot \begin{bmatrix} x \\ y \\ w \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{x_0} & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ w \end{bmatrix} = \begin{bmatrix} x \\ y \\ \frac{x}{x_0} + w \end{bmatrix}$$



# Field of view and viewing line



- All points on the affine Line  $x = -x_0$  are mapped to infinite points
- Only points on one side of this line are transformed
- These points are in the field of view, and the line  $x = -x_0$  is the viewing line





# Fixed points and lines

- Points on the projection line  $x=0$  (=  $y$ -axis) together with the infinite point  $[0, 1, 0]$  are fixed points of this transformation (the  $y$ -axis is invariant)
- Lines parallel to the  $y$ -axis remain parallel



# Fixed points and lines



- Points on the projection line  $x=0$  (=  $y$ -axis) together with the infinite point  $[0, 1, 0]$  are fixed points of this transformation (the  $y$ -axis is invariant)
- Lines parallel to the  $y$ -axis remain parallel

$$T_p \cdot \begin{bmatrix} x_{obj} \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ \frac{1}{x_0} & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} x_{obj} \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} x_{obj} \\ y \\ \frac{x_{obj}}{x_0} + 1 \end{bmatrix} = \begin{bmatrix} x_{obj} \\ y \\ \frac{x_{obj} + x_0}{x_0} \end{bmatrix} = \begin{bmatrix} \frac{x_{obj} \cdot x_0}{x_{obj} + x_0} \\ \frac{y \cdot x_0}{x_{obj} + x_0} \\ 1 \end{bmatrix} = \begin{bmatrix} x_{image} \\ y \cdot a_{factor} \\ 1 \end{bmatrix}$$

- Lines are transformed to lines



# Parallel lines

- The affine (eye) point  $[-x_0, 0, 1]^t$  is transformed to the infinite point  $[-x_0, 0, 0]^t = [-1, 0, 0]^t$
- The points on the affine  $y$ -axis are invariant



# Parallel lines

- Lines are mapped to lines
  - A line through eye point  $[-x_0, 0, 1]^t$ , which intersects the  $y$ -axis at  $[0, y_0, 1]^t$  is mapped to a line parallel to the  $x$ -axis passing through  $[0, y_0, 1]^t$





# Vanishing line

- The point  $[x, y, 0]^t$  is mapped to  $[x_0, x_0 \cdot y/x, 1]^t$ 
  - Note that  $[x, y, 0]^t$  is a **direction** and  $[x_0, x_0 \cdot y/x, 1]^t$  is a **point**
- The mappings of all lines parallel to the affine line with direction  $[x, y, 0]^t$  contain the point  $[x_0, x_0 y/x, 1]^t$ , meaning they all intersect in this point
- The union of the mappings of all lines with direction  $[x, y, 0]^t$  lie on the line  $x=x_0$  (= vanishing line)

# Vanishing line

- The point  $[x, y, 0]^t$  is mapped to  $[x_0, x_0 \cdot y/x, 1]^t$ 
  - Note that  $[x, y, 0]^t$  is a **direction** and  $[x_0, x_0 \cdot y/x, 1]^t$  is a **point**







# One, two and three vanishing point perspectives

- General perspective transformation

$$T_p \cdot \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ \frac{1}{x_0} & \frac{1}{y_0} & \frac{1}{z_0} & 1 \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} = \begin{bmatrix} x \\ y \\ z \\ \frac{x}{x_0} + \frac{y}{y_0} + \frac{z}{z_0} + w \end{bmatrix}$$

- The directions of lines parallel to the coordinate axes are mapped to the vanishing points  $[x_0, 0, 0, 0]^t$ ,  $[0, y_0, 0, 0]^t$ ,  $[0, 0, z_0, 0]^t$






# In OpenGL code

- For example, in `reshape (...)`  
`glViewport(0, 0, width, height)`  
`glMatrixMode(GL_PROJECTION)`  
`glLoadIdentity()`  
`gluPerspective(...)` ← **Perspective transformation**  
`glMatrixMode(GL_MODELVIEW)`
- In `render ()`  
`gluLookAt(...)` ← **Viewing transformation**  
`glTranslatef(...)`  
`glRotatef(...)` ← **Modeling transformations**  
`draw_scene ()`


# Geometric construction




# Geometric construction

This transformation maps points  $[x,y,0]$  to  $[x,y,x/4]$


The vanishing point of parallel lines AB and DC can be computed from the direction  $[4,1,0]$  as  $[4,1,1]$ , which is the 2D point  $[4,1]$



# Geometric construction




# Geometric construction






# Geometric construction



# Geometric construction





# Geometric construction

